

Experimental Investigations of Signals of a New Nature with the aid of two High Precision Stationary Quartz Gravimeters.

Yu.A.BAUROV*

*Central Research Institute of Machine Building
141070, Moscow region, Korolev, Pionerskaya, 4, Russia*

A.V.KOPAJEV

*Shternberg State Astronomical Institute of Moscow State University (SSAI MSU)
119899, Moscow, University prospect, 13, Russia*

In consequence of long-term (2000) observation of a system of two high-precision quartz gravimeters (one of them with an attached magnet) placed in a special (at a depth of ~ 10 m) gravimetric laboratory on a common base separated from the foundation of the building, signals of a new nature were detected. They were of a smooth peak-type shape, several minutes duration, and with amplitudes often more than that of the moon tide.

The nature of the signals cannot be explained in the framework of traditional physical views but can be qualitatively described with the aid of a supposed new interaction connected with the hypothesis about the existence of the cosmological vectorial potential $\mathbf{A}_g$, a new presumed fundamental vectorial constant.

PACS numbers: 04.80.+Z, 11.10.Lm, 98.80.-k

I. INTRODUCTION.

The signal of a new nature were first detected in a run of experiments of 1994-1996, carried out with the aid of a modernized high-precision quartz gravimeter "Sodin" and a magnet attached to it [1-4]. The signal appeared as permanent smooth peak-shape pulses with duration from 2 to 10min. and with amplitudes up to $\sim 15L$ (L is the amplitude of the moon tide). The experiments were carried out within a special box of the gravimetric laboratory of SSAI MSU, in a basement room at a level of 2 meters underground. The duration of the uninterrupted experiment was no more than 30 days. Besides the last peak (19.04.96), all results were obtained on the basis of one gravimeter measurements. In the last experiment two gravimeters were used. They were positioned on the same base in the box: one of the them was with a magnet, the other (reference one) was without magnet. Both gravimeters were connected to the same power network but had separated stabilized power sources.

In the experiment with two gravimeters, that with magnet detected an anomalous event which was not fixed by the gravimeter without magnet.

Since the procedure of investigating the signal of new nature with the aid of two gravimeters was of short-term duration (and the main results were obtained by one gravimeter), the run of experiments in 1994-1996 could be considered only as preliminary one. Nonetheless, it should be noted that the signals (12 peaks) were concentrated in space around some stellar direction having the following coordinates in the second equatorial system: right ascension $\alpha \approx 270^\circ$, declination $\delta \approx 34^\circ$. This was nearly coincident with some chosen spatial direction found in the course of investigating the assumed new anisotropic interaction with the aid of a torsion balance arranged inside high-current magnets [3-8] as well as when observing time changes in the intensity of the β -decay of radioactive elements [3,4,9-10] in the process of their rotation together with the Earth.

The new run of experiments was aimed at more precise measurements owing to elimination of technique errors of the preceding investigations.

II. THE EXPERIMENTAL SETUP AND PROCEDURE.

In the run of experiments of 1999-2000, the method of two gravimeters with a magnet attached to one of them, was used. The gravimeters were positioned in an underground gravimetric laboratory of SSAI MSU, at a depth of ~ 10 m, on a special base separated from the foundation of the building. A recently made stabilized power system could feed the gravimeters also in a case of accidental power switching off during ~ 2 hs. The information from gravimeters was transmitted to a personal computers (PC).

A schematic diagram of a quartz sensitive system is shown in Fig.1. Its main component is a quartz lever 1 (2 cm in length with a platinum mass of $m = 0.05g$) suspended on torsion quartz fibres 2 and additionally off-loaded by a vertical quartz spring 3. Such a construction gives to the device the high sensitivity to changes in gravity up to $1\mu Gal$ ($10^{-8}ms^{-2} = 10^{-9}G$, G is the free fall acceleration) as well as sufficient protection against microseismic disturbances.

The optical recording system comprises a galogen lamp 4 fed from a high-stabilised power source, light of which

*baurov@www.com

lamp enters into the instrument through the objective lens 5 with the aid of optical fibers. The quartz rod 6 welded to the level is a cylindrical lens, it forms an image of the point light source, which falls further on a photosensor rule 7 with a sensitivity on the order of $1\mu\text{m}$. The digital signal from the photosensor rule output enters further through a special interface to a computer PC which executes preliminary processing and averaging of data over the interval of 1 min. The accuracy of minute's data is $(0.5 - 1.2) \times 10^{-9}G$ depending on the microseismic noise level varying considerably in the course of day.

The sensitive system of the instrument includes some additional units (not shown in Fig.1) protecting it from thermal, electrostatic, and atmospheric pressure disturbances.

Calibration of the instrument is carried out by the inclination procedure with an accuracy of about $(0.5 - 1.0)\%$ which is quite sufficient for the experiment being considered.

The amplitude of constantly recorded by the device changes in gravity due to Moon-Sun tides and corresponding Earth's deformation, is $(50 - 200) \times 10^{-9}G$ which allows to evaluate magnitudes of possible anomalous effects against the background of tides.

To measure the new interaction by the Sodin gravimeter, a constant magnet (60 mm in diameter, 15mm in height, the field B in the centre of $0.3T$) was attached to it in such a way that the vector potential lines of the magnet in the vicinity of a test platinum weight (see Fig.1) were directed perpendicular to the Earth's surface. The magnet played the role of a peculiar amplifier of the new force (see in detail Refs. [3,4]).

III. RESULTS OF EXPERIMENTAL INVESTIGATIONS.

To illustrate the experiment, in Figs.2-6 some temporal fragments of recording of the gravimeters are shown. They demonstrate smooth peak-form overshoots by the "Sodin" gravimeter S209 with the attached constant magnet.

At the first stage of experiments, the gravimeter with the magnet and the reference gravimeter worked on different time scales (Moscow (MSK) and Greenwich (UTC) time, respectively). Therefore a shift of the curves is seen in Figs.2,3 which corresponds to that of time zones (i.e. 3hs).

It is seen from the Figures that the amplitudes of overshoots are in some cases (Figs.4,5) much superior to the amplitude of the moon tide. In February-March 2000, the overshoots corresponded to an increase in the gravitational effect, in May-June on the contrary to its decrease.

As opposed to the experiments of 1994-1996 in which the duration of the peaks was various (from 2 to 10 min. and more), the peak length in consideration was practically constant and equaled 2-3 min.

In Table 1 presented are all anomalous deflections recorded by the gravimeter with magnet and not fixed by the reference gravimeter without magnet.

Consider in detail an experimental fragment shown in Fig.6. As is seen, the events 1,3,4 corresponding to local earthquakes are neatly coincident in the recordings of both gravimeters, but the event 2 was fixed only by the gravimeter with magnet.

IV. DISCUSSION.

As was said above, the instruments in the course of the experiment were arranged on the base separated from the foundation of the building, at a depth of $\sim 10\text{m}$. Therefore, as is clearly seen from Fig.6, various Earth's oscillations acted identically onto both gravimeters.

Their design gives them good protection against thermal, electrostatic, and barometric disturbances. The gravimeters were powered from the same network, therefore, if some disturbance of "unknown nature"¹ could pass through it that would manifest itself in both readings. The possible "jumps" in quartz occurring sometimes when using the gravimeters "Sodin" have quite different form and cannot explain the nature of the smooth minute's peaks fixed in the experiment. The result obtained also cannot be explained by noise in the computers, because during the long use of them before the run of experiments of 1994-1996 as well as in other experiments carried out in 2001, such peaks were not observed. The outer electromagnetic noise was practically minimum and, which is the main, that would uniformly tell on both gravimeters.

One cannot also explain the obtained results by the influence on the gravimeter of various hypothetical particles predicted in modern calibration theories (like axions, photinos, gravitinos, etc.) [11,12] since both gravimeters are equal for their properties.

A temporal analysis of peak's appearance in readings of the gravimeter in association with the solar activity has shown the following (See Table 1).

In Table 1 shown are the times of detection of the peaks by the gravimeter with magnet during the first half-year 2000 as well as times of corresponding events in the Sun with indication of character of the event and the observatory from which the information was obtained. The more complete information about the solar events accompanying the peaks can be found in Refs. [13].

As is seen from Table 1, the peak detected 24.03.2000 by the gravimeter with magnet coincides practically precisely with a sweep-frequency radio burst.

¹ The electric power supply of the instrumentation was stabilized, so that there was no problem with it and it was not switched off.

The event (2) in Fig.6 occurred 5 minutes before a potent solar flare. The flares of such potency were not observed earlier during more than 40 years.

The other events fixed by the gravimeter lie several minutes or tens of minutes earlier or later than solar events of ordinary importance.

The results obtained can be qualitatively explained with the help of assumed new interaction caused by the existence of the vector $\mathbf{A}_g$.

According to recent investigations of anisotropic properties of the new force presented in Ref.[14], this force is directed at each point of space over a cone with the opening of $90 - 100^\circ$ formed around the vector $\mathbf{A}_g$ with the coordinates $\alpha \approx 293^\circ \pm 10^\circ$; $\delta \approx 36^\circ \pm 10^\circ$. The results of [14] are in good agreement with [15].

The force can act also in direction of $\mathbf{A}_g$ if an object being acted upon has a closed circular current. This was observed in experiments on investigating changes in the intensity of β -decay of radioactive elements [15].

An analysis of the experimental material presented here has shown that all events in Table 1 (except (2) in Fig.6) corresponded to such observation time when the direction of a normal to the surface of the Earth (i.e. the direction of the maximum sensitivity of the gravimeters) was coincident (to a precision of $\pm 10^\circ$) with the generator of above said cone. The time of the event (2) in Fig.6 indicates the direction of the normal to the Earth's surface coincident (to $\pm 5^\circ$) with the coordinates of the vector $\mathbf{A}_g$ itself: $\alpha = 293^\circ$, $\delta = 36^\circ$ (i.e. with the axis of the cone).

It was shown in Ref.[16] that the observed anisotropy of solar flares reveals a very similar direction ($\alpha = 277^\circ \pm 5^\circ$, $\delta = 38^\circ \pm 5^\circ$) which can be explained on the basis of the new anisotropic interaction.

Thus the experiments carried out with the use of two gravimeters have confirmed the earlier results obtained with the aid of only one gravimeter with an attached magnet. Their results (the direction of the new force) are coincident with those of the experiments on investigating the anisotropic properties of space with the aid of plasma generators as well as the changes in intensity of the β -decay of radioactive elements.

The authors are grateful to participants of the seminar at the Astrocsmic center of the Physical Institute of RAS and personally to prof. V.V.Burdyuzsa for a fruitful discussion of the experimental results, as well as to E.P.Morozov, L.I.Kazinova, A.Yu.Baurov for the help in preparation of the text of the paper.

- [4] Yu.A.Baurov, On the structure of physical vacuum and a new interaction in Nature (Theory, Experiment and Applications), Nova Science, NY, 2000.
- [5] Yu.A.Baurov, E.Yu.Klimenko, S.I.Novikov, Dokl. Acad. Nauk (DAN), **315**, 5, 1990, p.1116 (in Russian)
- [6] Yu.A.Baurov, P.M.Ryabov, Dokl. Acad. Nauk (DAN), **326**, N1, 1992, p.73 (in Russian).
- [7] Yu.A.Baurov, E.Yu.Klimenko, S.I.Novikov, Phys. Lett. **A162**, 1992, p.32.
- [8] Yu.A.Baurov, Phys.Lett. **A181**, 1993, p.283.
- [9] Yu.A.Baurov, V.L.Shutov, Applied Physics, **1**, 1995, p.40 (in Russian).
- [10] Yu.A.Baurov, Yu.G.Sobolev, V.F.Kushniruk, E.A.Kuznetsov and A.A.Konradov, E-print hep-ex/9907008.
- [11] H.V.Klapdor-Kleingrothaus, K.Zuber, Teilchenasto Physik, B.G.Teubner, GmbH. Stuttgart, 1997.
- [12] M.Green, J.H.Schwarz, E.Vitten, Theory of Superstrings, in 2 vol., Mir, Moscow, 1990 (in Russian).
- [13] <http://www.spaceweather.com>, <http://sec.noaa.gov>.
- [14] Yu.A.Baurov, I.B.Timofeev, V.A.Chernikov, S.F.Chalkin, E-print physics/0102090.
- [15] Yu.A.Baurov, G.A.Beda, I.P.Danilenko and V.P.Ignatko, Hadronic Jornal Supplement, **15**, (2000), 195.
- [16] Yu.A.Baurov, A.A.Konradov, V.F.Kushniruk, E.A.Kuznetsov, Yu.V.Ryabov, A.P.Senkevich, Yu.G.Sobolev, S.V.Zadorozsny, E-print hep-ex/0105009.
- [17] Yu.A.Baurov, A.A.Efimov, A.A.Shpitalnaya, E-print gr-qc/9606033.

N	S209 events	The events in the Sun				
		Begin	Max	End	Obs.	Type
1.	07.02.2000 09 ^h	08 ^h 54 ^m 09 ^h 42 ^m	/ / / / 09 ^h 50 ^m	08 ^h 55 ^m 09 ^h 57 ^m	LEA LEA	RSP FLA
2.	04.03.2000 06 ^h 05 ^m	05 ^h 41 ^m 06 ^h 20 ^m	05 ^h 42 ^m 06 ^h 28 ^m	05 ^h 57 ^m 07 ^h 08 ^m	LEA LEA	FLA FLA
3.	24.03.2000 01 ^h 08 ^m	00 ^h 35 ^m 01 ^h 08 ^m 03 ^h 09 ^m	00 ^h 40 ^m / / / / 03 ^h 14 ^m	00 ^h 47 ^m 01 ^h 09 ^m 03 ^h 17 ^m	GO8 LEA GO8	XRA RSP XRA
4.	26.05.2000 17 ^h 05 ^m	16 ^h 25 ^m 18 ^h 02 ^m	16 ^h 26 ^m 18 ^h 07 ^m	16 ^h 29 ^m 18 ^h 16 ^m	HOL GO8	FLA XRA
5.	27.05.2000 15 ^h 36 ^m	14 ^h 45 ^m 15 ^h 49 ^m	/ / / / 15 ^h 50 ^m	14 ^h 45 ^m 16 ^h 00 ^m	SVI RAM	RSP FLA
6.	06.06.2000 12 ^h 01 ^m	12 ^h 00 ^m 12 ^h 06 ^m 12 ^h 08 ^m	12 ^h 18 ^m 15 ^h 21 ^m 12 ^h 12 ^m	12 ^h 25 ^m 18 ^h 43 ^m 12 ^h 13 ^m	GO8 RAM SVI	XRA FLA RBR

Table 1. The events were recorded by the gravimeter with magnet and the events in the Sun.

Obs. (the reporting observatory): GO8 - GOES-8 satellite; HOL - Holloman AFB, NM, USA; SVI - San Vito, Italy; LEA - Learmonth, Australia; RAM - Ramey AFB, PR, USA.

Type (types of events): FLA - Optical flare observed in H-alpha; RBR - Fixed-frequency radio burst; RSP - Sweep-frequency radio burst; XRA - X-ray flare.

Begin, Max, End - the UTC time of the beginning, maximum, and the end of the event, / / / / - indicates a missing time.

[1] Yu.A.Baurov, A.V.Kopajev, LANL E-print hep-ph/9601369.

[2] Yu.A.Baurov, A.V.Kopajev, Fizicheskaya Mysl Rossii, **2**, 1996, 1 (in Russian).

[3] Yu.A.Baurov, Structure of Physical Space and New Method of Generating Energy (theory, experiment, applied issues), Moscow, "Krechet", 1998 (in Russian).




Fig.1 Sensitive system of the Sodin quartz gravimeter:
 1 - beam,
 2 - horizontal quartz wires suspension system,
 3 - main spring,
 4 - lamp,
 5 - object lens,
 6 - beam,
 7 - CCD-SCALE,
 8 - prism,
 9, 10 - thermocompensator,
 11, 12, 13, 14 - micrometric compensation mechanism,
 15 - constant magnet.




Fig.2 Readings of the gravimeters at Mar.04.2000.
 y - the displacement of platinum weight (one division is equal to 0.1μ and corresponds to $0.2\mu Gal$);
 x - time (in minutes);
 Event (peak) at $6^h 5^m$ UTC ($9^h 5^m$ MSK).




Fig.3 The same as in Fig.2 at Mar.24.2000.
Event (peak) at $1^{h}8^{m}$ UTC ($4^{h}8^{m}$ MSK).




Fig.4 The same as in Fig.2 at May.26.2000.
Event (peak) at $17^{h}5^{m}$ UTC.




Fig.5 The same as in Fig.2 at May.27.2000.
Event (peak) at 15^h36^m UTC.




Fig.6 The same as in Fig.2 at Jun.6.2000.
Event (peak) at 12^h1^m UTC.